

大分大学 福祉健康科学部 理学療法コース 業績目録 2017

1. Anan M, Hattori H, Tanimoto K, Wakimoto Y, Ibara T, Kito N, Shinkoda K.
The coordination of joint movements during sit-to-stand motion in old adults: the uncontrolled manifold analysis.
Phys Ther Res. 2017;**20**(2):44–50. doi: [10.1298/ptr.E9923](https://doi.org/10.1298/ptr.E9923).
2. Sugata H, Hirata M, Tamura Y, Onishi H, Goto T, Araki T, Yorifuji S.
Frequency-dependent oscillatory neural profiles during imitation.
Sci Rep. 2017;**7**:45806. doi: [10.1038/srep45806](https://doi.org/10.1038/srep45806).
3. Itoh Y, Murakami T, Mori T, Agata N, Kimura N, Inoue-Miyazu M, Hayakawa K, Hirano T, Sokabe M, Kawakami K*.
Training at non-damaging intensities facilitates recovery from muscle atrophy.
Muscle Nerve. 2017;**55**(2):243–253. doi: [10.1002/mus.25218](https://doi.org/10.1002/mus.25218).
4. Shuto M, Fujioka T, Matsunari O, Okamoto K, Mizukami K, Okimoto T, Kodama M, Takigami S, Seguchi C, Nonaka Y, Sato R, Yamaoka Y, Murakami K.
Association between Gastric Cancer Risk and Serum Helicobacter pylori Antibody Titers.
Gastroenterol Res Pract. 2017;**2017**:1286198. doi: [10.1155/2017/1286198](https://doi.org/10.1155/2017/1286198).
5. Mizukami K, Ogawa R, Okamoto K, Shuto M, Fukuda K, Sonoda A, Matsunari O, Hirashita Y, Okimoto T, Kodama M, Murakami K.
Objective Endoscopic Analysis with Linked Color Imaging regarding Gastric Mucosal Atrophy: A Pilot Study.
Gastroenterol Res Pract. 2017;**2017**:5054237. doi: [10.1155/2017/5054237](https://doi.org/10.1155/2017/5054237).
6. Bouta EM, Kuzin I, de Mesy Bentley K, Wood RW, Rahimi H, Ji RC, Ritchlin CT, Bottaro A, Xing L, Schwarz EM.
Brief Report: Treatment of Tumor Necrosis Factor-Transgenic Mice With Anti-Tumor Necrosis Factor Restores Lymphatic Contractions, Repairs Lymphatic Vessels, and May Increase Monocyte/Macrophage Egress.
Arthritis Rheumatol. 2017;**69**(6):1187–1193. doi: [10.1002/art.40047](https://doi.org/10.1002/art.40047).
7. Tokuda K, Anan M, Sawada T, Tanimoto K, Takeda T, Ogata Y, Takahashi M, Kito N, Shinkoda K.
Trunk lean gait decreases multi-segmental coordination in the vertical direction.

J Phys Ther Sci. 2017;**29**(11):1940–1946. doi: [10.1589/jpts.29.1940](https://doi.org/10.1589/jpts.29.1940).

8. Tanimoto K, Takahashi M, Tokuda K, Sawada T, Anan M, Shinkoda K.
Lower limb kinematics during the swing phase in patients with knee osteoarthritis measured using an inertial sensor.
Gait Posture. 2017;**57**:236–240. doi: [10.1016/j.gaitpost.2017.06.017](https://doi.org/10.1016/j.gaitpost.2017.06.017).
9. Sawada T, Tanimoto K, Tokuda K, Iwamoto Y, Ogata Y, Anan M, Takahashi M, Kito N, Shinkoda K.
Rear foot kinematics when wearing lateral wedge insoles and foot alignment influence the effect of knee adduction moment for medial knee osteoarthritis.
Gait Posture. 2017;**57**:177–181. doi: [10.1016/j.gaitpost.2017.06.009](https://doi.org/10.1016/j.gaitpost.2017.06.009).
10. Hashimoto H, Hasegawa Y, Araki T, Sugata H, Yanagisawa T, Yorifuji S, Hirata M.
Non-invasive detection of language-related prefrontal high gamma band activity with beamforming MEG.
Sci Rep. 2017;**7**(1):14262. doi: [10.1038/s41598-017-14452-3](https://doi.org/10.1038/s41598-017-14452-3).

* corresponding author